



Wlodarczyk, A. I., Sylantyev, S., Herd, M. B., Kersanté, F., Lambert, J. J., Rusakov, D. A., Linthorst, A. C. E., Semyanov, A., Belelli, D., Pavlov, I., & Walker, M. C. (2013). GABA-independent GABAA receptor openings maintain tonic currents. *Journal of Neuroscience*, 33(9), 3905-3914. <https://doi.org/10.1523/JNEUROSCI.4193-12.2013>

Publisher's PDF, also known as Version of record

Link to published version (if available):
[10.1523/JNEUROSCI.4193-12.2013](https://doi.org/10.1523/JNEUROSCI.4193-12.2013)

[Link to publication record in Explore Bristol Research](#)
PDF-document

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

GABA-Independent GABA_A Receptor Openings Maintain Tonic Currents

Agnieszka I. Włodarczyk,^{1*} Sergiy Sylantyev,^{1*} Murray B. Herd,² Flavie Kersanté,³ Jeremy J. Lambert,² Dmitri A. Rusakov,¹ Astrid C. E. Linthorst,³ Alexey Semyanov,^{4,5} Delia Belelli,² Ivan Pavlov,¹ and Matthew C. Walker¹

¹UCL Institute of Neurology, London WC1N 3BG, United Kingdom, ²Division of Neuroscience, Medical Research Institute, Ninewells Hospital & Medical School, Dundee University, Dundee DD1 9SY, United Kingdom, ³Henry Wellcome Laboratories for Integrative Neuroscience and Endocrinology, School of Clinical Sciences, University of Bristol, Bristol BS2 8AE, United Kingdom, ⁴RIKEN Brain Science Institute, Wako-shi 351-0114, Japan, and ⁵Nizhny Novgorod State University, Nizhny Novgorod 603000, Russia

Activation of GABA_A receptors (GABA_ARs) produces two forms of inhibition: phasic inhibition generated by the rapid, transient activation of synaptic GABA_ARs by presynaptic GABA release, and tonic inhibition generated by the persistent activation of perisynaptic or extrasynaptic GABA_ARs, which can detect extracellular GABA. Such tonic GABA_AR-mediated currents are particularly evident in dentate granule cells in which they play a major role in regulating cell excitability. Here we show that in rat dentate granule cells in *ex vivo* hippocampal slices, tonic currents are predominantly generated by GABA-independent GABA_A receptor openings. This tonic GABA_AR conductance is resistant to the competitive GABA_AR antagonist SR95531 (gabazine), which at high concentrations acts as a partial agonist, but can be blocked by an open channel blocker, picrotoxin. When slices are perfused with 200 nM GABA, a concentration that is comparable to CSF concentrations but is twice that measured by us in the hippocampus *in vivo* using zero-net-flux microdialysis, negligible GABA is detected by dentate granule cells. Spontaneously opening GABA_ARs, therefore, maintain dentate granule cell tonic currents in the face of low extracellular GABA concentrations.

Introduction

In addition to fast synaptic GABA_A receptor (GABA_AR)-mediated signaling, there is a slower form of signaling resulting from the tonic activation of GABA_ARs (Semyanov et al., 2004; Farrant and Nusser, 2005; Glykys and Mody, 2007a). Tonically active GABA_ARs can have profound effects on neuronal excitability, synaptic plasticity, network oscillations, and neurogenesis (Ge et al., 2006; Pavlov et al., 2009; Holter et al., 2010; Mann and Mody, 2010; Martin et al., 2010; Dubeau et al., 2011) and have been implicated in neuronal development, information processing, cognition, and memory (Semyanov et al., 2004; Farrant and Nusser, 2005; Brickley and Mody, 2012). The conventional view is that tonic currents are mediated by high-affinity extrasynaptic GABA_ARs that can detect low concentrations of ambient GABA, and therefore the magnitude of tonic currents is regulated by the expression of these receptors and the availability of extracellular

GABA (Semyanov et al., 2004; Farrant and Nusser, 2005; Glykys and Mody, 2007a).

The sources and concentration of extracellular GABA ([GABA]_e) are, however, still debated. Vesicular release, “leak” through bestrophin channels, and reversal of GABA transporters have all been suggested to contribute to the extracellular GABA pool (Attwell et al., 1993; Gaspary et al., 1998; Lee et al., 2010). *In vivo* estimates indicate that [GABA]_e is in the micromolar or submicromolar range (Lerma et al., 1986; Kuntz et al., 2004; Nyitrai et al., 2006). *In vitro* studies have suggested even lower concentrations of GABA, in the nanomolar range; indeed, active uptake maintains GABA at sufficiently low concentrations to prevent tonic GABA_B receptor activation (Isaacson et al., 1993). Thus, tonic currents in hippocampal neurons have often been measured in conditions that artificially raise [GABA]_e by inhibiting GABA uptake or metabolism, or by adding exogenous GABA to the perfusate (Overstreet and Westbrook, 2001; Nusser and Mody, 2002; Stell and Mody, 2002; Holter et al., 2010). When GABA is not increased by these means, various magnitudes of tonic currents have been obtained using different GABA_AR antagonists (Bai et al., 2001; Mtchedlishvili and Kapur, 2006; Zhan and Nadler, 2009; Mtchedlishvili et al., 2010). Here we report a novel form of tonic inhibition mediated by GABA-independent openings of GABA_ARs, which can explain these discrepancies.

We focus on dentate gyrus granule cells (DGCs), the excitability of which critically depends upon the presence of tonic GABA_AR-mediated conductances (Overstreet and Westbrook, 2001; Nusser and Mody, 2002; Stell and Mody, 2002; Coulter and Carlson, 2007; Holter et al., 2010). We demonstrate that under

Received Aug. 31, 2012; revised Jan. 2, 2013; accepted Jan. 12, 2013.

Author contributions: A.I.W., S.S., M.B.H., F.K., J.J.L., D.A.R., A.C.E.L., A.S., D.B., I.P., and M.C.W. designed research; A.I.W., S.S., M.B.H., F.K., and I.P. performed research; A.I.W., S.S., M.B.H., F.K., J.J.L., D.A.R., A.C.E.L., A.S., D.B., I.P., and M.C.W. analyzed data; A.I.W., I.P., and M.C.W. wrote the paper.

*A.I.W. and S.S. contributed equally to this work.

This work was supported by Wellcome Trust Grants WT083163 (A.I.W., A.S., F.K., A.C.E.L., and M.C.W.) and WT084311 (D.A.R.); Medical Research Council Grants G1000008 (D.B. and J.J.L.), and G0900613 and G0802216 (D.A.R.); Tenovus Scotland (J.J.L. and D.B.); Anonymous Trust (J.J.L. and D.B.); Epilepsy Research UK Grants F1001 (M.B.H.) and A0832 (I.P.); and Worshipful Company of Pewterers (I.P.).

Correspondence should be addressed to either Ivan Pavlov or Matthew C. Walker, Department of Clinical and Experimental Epilepsy, UCL Institute of Neurology, London WC1N3BG, UK. E-mail: i.pavlov@ucl.ac.uk or mwalker@ion.ucl.ac.uk.

DOI:10.1523/JNEUROSCI.4193-12.2013

Copyright © 2013 the authors 0270-6474/13/333905-10\$15.00/0

baseline conditions, or when the perfusate contains the same concentration of GABA that is found in CSF, the major contributors to the tonic current in DGCs are spontaneously opening GABA_ARs. This tonic GABA_AR conductance is resistant to the competitive GABA_AR antagonist gabazine (SR95531), but can be blocked by an open channel blocker, picrotoxin. On increasing [GABA]_e, a SR95531-sensitive component of tonic current emerges. Together, these results indicate that the GABA-independent component of tonic current mediated by spontaneously opening GABA_ARs maintains tonic inhibition in the presence of low extracellular GABA concentrations measured *in vivo*.

Materials and Methods

Hippocampal slice preparation

Transverse hippocampal slices (350 μm thick) were used for *in vitro* electrophysiological recordings. Slices were prepared from 3- to 4-week-old male Sprague Dawley rats and δ^{-/-} knockouts or δ^{+/-} littermate control mice on a C57B6 background (Mihalek et al., 1999; Herd et al., 2008). Animals were killed by an overdose of isoflurane according to the United Kingdom Animals (Scientific Procedures) Act of 1986. After decapitation, brains were rapidly removed and dissected, and hippocampi were sliced with a Leica VT1200S vibratome in ice-cold sucrose-based solution containing the following (in mM): 70 sucrose, 80 NaCl, 2.5 KCl, 7 MgCl₂, 0.5 CaCl₂, 25 NaHCO₃, 1.25 NaH₂PO₄, 22 glucose, equilibrated with 95% O₂ plus 5% CO₂, pH 7.4, 315–330 mOsm. Slices were maintained in continuously oxygenated sucrose-free storage solution at 33°C for 15 min, equilibrated to a room temperature for 15 min and then placed to recover in continuously oxygenated humid interface holding chamber at room temperature for at least 1 h before recording. In experiments with concanamycin, slices were prepared in exactly the same way, but before placing them in a holding chamber for the recovery they were incubated for 2 h in a submerged chamber with continuously oxygenated storage solution and 0.5 μM concanamycin. After that, slices were placed in an interface holding chamber and allowed to rest for about half an hour before recording. After recovering slices were transferred into recording chamber. The perfusion and storage medium contained the following (in mM): 119 NaCl, 2.5 KCl, 1.3 MgSO₄, 2.5 CaCl₂, 26.2 NaHCO₃, 1 NaH₂PO₄, 22 glucose and was gassed with 95% O₂ and 5% CO₂, pH 7.4; 290–298 mOsm.

In vitro electrophysiology

Whole-cell recordings. Visualized patch-clamp recordings from mature dentate granule cells (input resistance, R_{in} = 310 ± 50 MΩ; membrane capacitance, C_m = 48 ± 6 pF) were performed using an infrared differential interference contrast imaging system. Tonic GABA_AR-mediated currents were measured in voltage-clamp mode (at holding potential V_{hold} = −70 mV) in the presence of ionotropic glutamate receptor blockers DL-APV (50 μM) and NBQX (20 μM), metabotropic glutamate receptors blocker (S)-α-methyl-4-carboxyphenylglycine (MCPG; 250 μM), and GABA_B receptor blocker (2S)-3-[[[(1S)-1-(3,4-dichlorophenyl)ethyl]amino-2-hydroxypropyl](phenylmethyl)phosphinic acid (CGP55845; 1 μM). The intracellular pipette solution contained the following (mM): 120.5 CsCl, 10 KOH-HEPES, 2 EGTA, 8 NaCl, 5 QX-314 Br[−] salt, 2 Mg-ATP, 0.3 Na-GTP, 10 Na-phosphocreatine, pH adjusted to 7.2 and osmolarity adjusted to 295 mOsm. The tonic GABA_AR-mediated current was measured as an outward shift in holding current following application of picrotoxin (100 μM). Changes in root mean square (RMS) noise have also been proposed to reflect changes in tonic GABA_AR-mediated conductances and have been used because they are unaffected by current drift. Although RMS noise decreased in experiments in which tonic currents were blocked, this measurement is confounded by the presence of synaptic currents. Moreover, RMS noise is non-linearly related to current and can paradoxically decrease when tonic currents increase (Traynelis and Jaramillo, 1998; Glykys and Mody, 2007a). We, therefore, only used RMS noise as a measure in experiments in which we were trying to block the tonic current. Three other GABA_AR antagonists, bicuculline methiodide (BMI) (10 μM), SR95531 (0.5, 25, and 125 μM), and pentylenetetrazol (1.5 mM) were used in experiments. Gabazine (25 μM) and

bicuculline (10 μM) were used at concentrations that would be predicted to give >90% occupancy of GABA_ARs in dentate granule cells (Jones et al., 2001). The affinity of gabazine for GABA_ARs in dentate granule cells is an order of magnitude greater than that of bicuculline (Jones et al., 2001), so that equimolar concentrations of gabazine displace bicuculline from its binding site. Glycine receptors were blocked with 1 μM strychnine, GABA_C receptors—with 50 μM (1,2,5,6-tetrahydropyridin-4-yl)methylphosphinic acid (TPMPA). Recordings were performed at 32–34°C. The whole-cell pipette resistance was 4–5.5 MΩ. Series resistance was monitored throughout experiments using a −5 mV step command. Cells showing a >20% change in a series resistance, a resistance >20 MΩ, or unstable holding current were rejected.

Acquisition and analysis. Recordings were obtained using a MultiClamp 700B or Axopatch 200B amplifier (Molecular Devices), filtered at 4 kHz, digitized at 10 kHz, and stored on a PC. WinEDR (Strathclyde Electrophysiology Software) was used for data acquisition and pClamp (Molecular Devices), and OriginPro 8.1 (OriginLab) for off-line analysis.

For analysis of tonic currents, mean values of holding current during 200 ms epochs free of synaptic currents were measured every 30 s. The amplitude of the tonic current was calculated as the difference between the holding current (ΔI_{hold}) measured 6 min before and 10 min after the application of an antagonist. The speed of perfusion in our system of 3.5 ml/min allowed full equilibration of the solution in the recording chamber at this time point. The values of RMS noise were calculated for 200 ms epochs free of synaptic events. The change of RMS noise (Δrms) was calculated as the difference between the values before and after the application of an antagonist.

Outside-out and nucleated patch recordings. GABA_AR-mediated currents were recorded in the presence of 0.1 μM CGP55845, 100 μM D-APV, 10 μM NBQX, 200 μM S-MCPG, and 1 μM strychnine. The intracellular pipette solution contained the following (in mM): 120.5 CsCl, 10 KOH-HEPES, 10 BAPTA, 8 NaCl, 5 QX-314 Br[−] salt, 2 Mg-ATP, 0.3 Na-GTP, pH 7.2, osmolarity 295 mOsm. Patches were pulled from dentate granule cells, and recordings were performed in voltage-clamp mode at 33–34°C (V_{hold} = −70 mV) using MultiClamp 700B amplifier. Signals were digitized at 10 kHz. The patch pipette resistance was 5–7 MΩ.

For the rapid solution exchange we used a θ-glass application pipette with ~200 μm tip diameter attached to the micromanipulator. The position of the pipette was controlled by piezoelectric element (the speed of switch was 50–100 μs). One pipette channel was filled with the bath aCSF solution or bath solution plus 10 μM GABA; another channel had 10 μM GABA, 25 or 125 μM SR95531, 10 μM bicuculline, or 20 μM picrotoxin. Pressure was regulated by a PDES-02DX pneumatic micro ejector (npi) using compressed nitrogen separately in each of two channels. Solutions with SR95531, bicuculline, and picrotoxin were exchanged in a pipette channel (7–12 s) during the exposure of nucleated patch to the bath solution channel (Bennett and Kearns, 2000; Sylantyev et al., 2008).

Analysis of the single-channel recordings

All analyses were performed on stretches of data that were longer than 3 min. The opening frequency of GABA_AR-mediated channels was calculated as $N/\Delta t$, where N is the number of openings and Δt is the time of recording. N was counted using a detection threshold of 1.5 pA more negative than mean baseline and a minimum opening time of 0.2 ms. To calculate single-channel conductance and average open time, we first built all-points amplitude histograms and fitted them with a two-Gaussian function:

$$F = \frac{p_1 e^{-\frac{(n-m_1)^2}{2\sigma_1^2}}}{\sigma_1 \sqrt{2\pi}} + \frac{p_2 e^{-\frac{(n-m_2)^2}{2\sigma_2^2}}}{\sigma_2 \sqrt{2\pi}},$$

where m_1 and m_2 are the mode values of Gaussians, σ_1 and σ_2 are the SDs of corresponding modes, n is the value of electrical current, and p_1 and p_2 are the fitting constants. The channel conductance was calculated as $G = (m_2 - m_1)/V_m$.

The Newton–Raphson iteration method was used to obtain the value of n_{min} , which is the value of n at the minimum point of the function F in the interval $m_1 < n < m_2$. This was taken as the point of channel closure.

Since the signal was digitized at 10 kHz (i.e., at intervals of 0.1 ms), the average open time (in milliseconds) was calculated as follows (the factor of 10 in the denominator is to account for the 0.1 ms interval):

$$T = \frac{\sum_{n=n_{\min}}^{n_{\max}} [F(n)]}{10 \times N},$$

where $n_{\max} = 2m_2 - n_{\min}$.

All calculations were performed using the Wolfram Mathematica 6.2 software; the general algorithm of histograms construction and interpretation was adapted from Bennett and Kearns (2000). We compared results obtained using this method against those obtained by averaging all the threshold-detected openings for the data shown in Figure 3. There was no difference between these two methods (Pearson's correlation coefficient of 0.992 for open time, and 0.979 for conductance).

Drugs and reagents

GABA receptor antagonists and concanamycin were purchased from Tocris Bioscience. Other reagents were from either Ascent Scientific or Sigma-Aldrich.

In vivo microdialysis

Animals. Male Sprague Dawley rats (Harlan) were housed three per cage under standard housing conditions (lights on between 5:00 A.M. and 7:00 P.M.; 21–22°C; 50–60% relative humidity) with free access to food pellets and drinking water. Animals were handled daily (~5 min/rat) starting 1 week before surgery and continuing until the day of the insertion of the microdialysis probe. At the time of surgery, rats weighed ~240 g. Surgical procedures were performed under isoflurane (Merial Animal Health) anesthesia. Carprofen (Rimadyl, 4 mg/kg, s.c.; Pfizer) was administered for postoperative pain relief. All procedures were conducted in accordance with the United Kingdom's Animals (Scientific Procedures) Act 1986, and all efforts were made to minimize animal numbers and suffering.

Surgical and microdialysis procedures. Nine days before the start of the experiment, rats were surgically prepared for microdialysis of the hippocampus by stereotaxic implantation of a guide cannula (MAB 6.14.IC, Microbiotech/se AB) essentially as described in detail before (Droste et al., 2008). After surgery, rats were housed individually in Plexiglas cages (length × width × height = 27 × 27 × 35 cm) under similar housing conditions as described above.

Seven days after surgery, a microdialysis probe (polyethersulfone membrane, length 4 mm, 15 kDa cutoff, and outer diameter 0.6 mm; MAB 6.14.4, Microbiotech/se AB) was inserted via the guide cannula into the CA3–dentate gyrus region of the hippocampus (Linthorst et al., 1994) under short-lasting isoflurane anesthesia (see Fig. 3C, schematics). Rats were connected to a liquid swivel and a counterbalance arm (Microbiotech/se AB), allowing free movement in all directions. Fluorethylene polymer tubing with a dead volume of 1.2 μl per 100 mm length (Microbiotech) was used for all connections. Dead volumes were accounted for during the experiment. Microdialysis probes were perfused with sterile, pyrogen-free Ringer solution (Delta Pharma) at a flow rate of 2 μl/min using a micro-infusion pump (KDS220, KD Scientific).

The zero-net-flux experiment was started at 9:00 A.M. on the second day after the insertion of the microdialysis probe. Ten minute samples were collected throughout the complete experiment. After 1 h of baseline sampling, microdialysis probes were subsequently perfused with increasing concentrations of GABA (10, 20, 40, 80, 160 and 240 nM; Sigma-Aldrich) dissolved in Ringer solution (reverse microdialysis). Each GABA perfusion lasted 40 min during which four 10 min samples were collected. Microdialysis samples were collected using an automated, refrigerated fraction collector (CMA470, CMA Microdialysis AB) and were stored at –80°C for later HPLC analysis.

Histology. At the end of the experiment, rats were killed using an overdose of pentobarbital (Euthatal, 200 mg/kg body weight, i.p.; Merial Animal Health), and the brains were removed and stored in a 4% buffered paraformaldehyde solution. Histological examination was performed as described previously (Linthorst et al., 1994; Droste et al.,

2008). Only data from rats with correctly placed microdialysis probes were included in the analyses.

Measurement of GABA. HPLC with electrochemical detection was used to measure GABA in the microdialysates essentially as described previously with small modifications (de Groote and Linthorst, 2007). Briefly, GABA was separated on a TARGA C18 10 cm × 1 mm column (particle size 3 μm; Higgins Analytical) using filtered and degassed mobile phase (18% methanol, 0.1 M NaH₂PO₄, 0.2 mM EDTA, pH 4.72) pumped at 50 μl/min using an Alexys LC-100 pump (Antec Leyden BV). Standards and samples were injected onto the column using a thermostatically controlled (8°C) Alexys AS-100 autosampler (Antec Leyden BV). Before injection, 13 μl of standard or sample was derivatized with 2 μl of *o*-phthalaldehyde (OPA)/sulfite solution (1.6 mM OPA, 0.5% methanol, 1.88 mM Na₂SO₃, 11.25 mM Na₂B₄O₇) for 4 min to make GABA electrochemically active. Next, 10 μl of the derivatized mixture was injected onto the column and GABA was detected using a VT-03 electrochemical flow cell (Antec Leyden BV) set at +850 mV against an Ag/AgCl reference electrode. Both column and detector were housed in a Faraday-shielded oven (DECADE II, Antec Leyden BV) thermostatically controlled at 38°C. Chromatograms were recorded and analyzed using Alexys chromatography software (Antec Leyden BV). The detection limit for GABA at a signal-to-noise ratio of 3:1 was 11–15 fmol per injection on column.

Calculations. Perfusion of each concentration of GABA (C_{in} ; nM) was performed for 40 min. However, to ensure that dialysis across the membrane had reached a steady state, the first 10 min sample was discarded and only the three subsequent samples were used for the calculations described below.

After measurement of the concentration of GABA in the collected samples (C_{out} ; in nM), the difference between C_{out} and C_{in} (i.e., the net loss or gain of GABA in the dialysate) was calculated ($C_{out} - C_{in}$; in nM) for each concentration of GABA perfused. The baseline condition represented $C_{in} = 0$. Next, for each individual animal, $C_{out} - C_{in}$ was plotted against C_{in} and, after regression analysis (GraphPad 5.0), the concentration of zero-net-flux was determined as the concentration of C_{in} at which $C_{out} - C_{in} = 0$. Next, the mean zero-net-flux concentration (\pm SEM; $n = 5$) was calculated. For graphical purposes, the mean \pm SEM values for $C_{out} - C_{in}$ were calculated for each concentration of GABA perfused and were plotted against C_{in} (see Fig. 3D).

Statistics

Statistical comparisons were made using paired and unpaired (as indicated in the text/figure legends) Student's *t* test, and Wilcoxon signed ranks test (Fig. 1C, experiments). Differences were considered significant at $p < 0.05$. Data are presented in the text and figures as the mean \pm SEM.

Results

Tonic activation of GABA_ARs in DGCs does not require synaptically released GABA

Since the accumulation of GABA in the extracellular space may result from synaptic release, we asked whether depleting vesicular GABA by incubating hippocampal slices in the vesicular H⁺-ATPase inhibitor concanamycin (0.5 μM) would affect tonic currents. We confirmed that concanamycin abolished synaptic currents (Rossi et al., 2003); however, it had no significant effect on the magnitude of the tonic current revealed by application of the GABA_A antagonist picrotoxin. Application of picrotoxin produced a 11.9 ± 1.5 pA outward shift in the I_{hold} in control experiments, and a 13.3 ± 3.0 pA shift when slices were preincubated in concanamycin (Fig. 1A). These results indicate that in acute hippocampal slices tonic currents in DGCs are not dependent on the concentration of GABA in vesicles or on GABA release into the synapse under baseline conditions. Glycine receptors can contribute to inhibition in the hippocampus and may also be blocked by picrotoxin in a use-dependent manner (Danglot et al., 2004; Yang et al., 2007). We tested their contribution by adding the glycine receptor antagonist strychnine (1 μM) and observed that it had no effect on I_{hold} and did





Figure 1. Pharmacology of the GABA_A-mediated tonic currents in DGCs. **A**, Concanamycin (0.5 μM) blocks exocytosis but does not affect I_{tonic} . No significant difference between the change in I_{hold} or RMS noise in control (left trace, $n = 9$) and concanamycin-treated slices (right trace, $n = 6$) upon application of picrotoxin (PTX; 100 μM). **B**, I_{tonic} in DGCs is insensitive to SR95531 and abolished by picrotoxin. SR95531 (0.5 μM) partially inhibited sIPSCs, while high concentrations (125 μM) completely abolished sIPSCs. Picrotoxin induced an outward shift in I_{hold} ($p = 0.0049$; $n = 5$) and decreased RMS noise ($p = 0.0012$; $n = 5$). **C**, In the presence of GABA (5 μM), application of SR95531 (0.5 μM) reduced sIPSCs, induced an outward shift in I_{hold} ($p = 0.016$), and decreased RMS noise ($p = 0.008$; $n = 8$). SR95531 (25 μM) abolished sIPSCs, induced a further nonsignificant shift in I_{hold} ($p = 0.44$), and RMS noise ($p = 0.06$; $n = 5$). SR95531 (125 μM) affected neither I_{hold} nor RMS noise. Subsequent application of picrotoxin produced an outward shift in I_{hold} and reduced RMS noise ($p = 0.031$ and $p = 0.003$ respectively; $n = 6$). **D**, BMI (10 μM) blocked all synaptic activity, induced an outward shift in I_{hold} and reduced RMS noise ($p = 0.00019$ and $p = 0.0014$ respectively; $n = 6$). SR95531 (25 μM) resulted in inward current and an increase in RMS noise ($p = 0.0013$ and $p = 0.008$ respectively; $n = 6$), which were reversed by picrotoxin (100 μM) ($p = 0.0019$ and $p = 0.0048$ respectively; $n = 5$). Statistical comparisons were made using paired Student's *t* test in **A**, **B**, and **D**, and Wilcoxon signed ranks test in **C**. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$. ΔI_{hold} and ΔRMS noise values represent changes from the previous drug application.

not occlude the effect of picrotoxin (ΔI_{hold} following application of strychnine: -0.8 ± 0.8 pA, $n = 5$, $p = 0.40$ compared with control; consecutive application of picrotoxin: 15.6 ± 3.3 pA, $n = 4$, $p = 0.018$ compared with strychnine). We further ruled out the potential contribution of GABA_CRs to picrotoxin-induced shift in I_{hold} by applying the GABA_C antagonist TPMPA (50 μM), which also did not

affect the amplitude of the picrotoxin-sensitive tonic current (ΔI_{hold} following application of TPMPA: -0.14 ± 1.78 pA, $p = 0.9$ compared with control; picrotoxin: 11.7 ± 1.7 pA, $p = 0.006$ compared with TPMPA; $n = 4$). Thus, we conclude that neither glycine receptors nor GABA_CRs contribute to tonic conductance revealed by picrotoxin.

Table 1. The effect of different GABA antagonists in control conditions with no GABA added

Antagonist	GABA tonic current (ΔI_{hold} , pA)
PTX (100 μM)	11.9 ± 1.5 , $n = 9$, $p = 0.00004$ compared to control
Pentylenetetrazol (1.5 mM)	10.6 ± 3.6 , $n = 5$, $p = 0.04$ compared to control
Bicuculline (10 μM)	5.9 ± 0.6 , $n = 6$, $p = 0.0002$ compared to control
SR95531 (125 μM)	-6.7 ± 1.6 , $n = 5$, $p = 0.01$ compared to control
PTX (100 μM) in SR95531 (125 μM)	18.9 ± 3.4 , $n = 5$, $p = 0.005$ compared to SR95531 (125 μM)

**Figure 2.** δ -Subunit-containing GABA_A receptors are involved in generating picrotoxin-sensitive tonic currents in DGCs. The effects of SR95531 (25 μM) and picrotoxin (PTX, 100 μM) on holding currents recorded in wild-type (top trace) and knock-out mice lacking δ -subunit-containing GABA_ARs (bottom trace). Picrotoxin-sensitive tonic currents are reduced in the knock-out mice compared with the wild-type littermate controls ($n = 9$ and $n = 10$ for the $\delta^{+/+}$ and $\delta^{-/-}$ mice, respectively; ** $p < 0.01$, Student's unpaired t test). Histogram shows mean values \pm SEM.

Tonic currents in DGCs are insensitive to SR95531 under baseline conditions

These results suggest either that $[\text{GABA}]_e$ is maintained at a constant concentration despite decreased vesicular GABA release (e.g., by nonvesicular release) and/or that there is another mechanism involved in generating tonic currents that is independent of $[\text{GABA}]_e$.

To distinguish between these possibilities, we took advantage of the distinct pharmacologies of different GABA_A antagonists. Picrotoxin is an open channel blocker and acts as a noncompetitive GABA_A antagonist that has equivalent efficacy in blocking low-affinity synaptic and high-affinity extrasynaptic GABA_ARs (Stell and Mody, 2002). In contrast, SR95531 is a competitive GABA_A antagonist and displaces GABA from its binding site (Hamann et al., 1988). While low concentrations of SR95531 (0.5 μM) partially inhibited and high concentrations (125 μM) totally blocked sIPSCs, there was no significant increase in I_{hold} , which became more negative at high SR95531 concentrations (ΔI_{hold} , 6.7 ± 1.6 pA, $n = 5$, $p = 0.014$ compared with 0.5 μM SR95531; Fig. 1B; Table 1), suggesting a partial agonist effect (this effect was not observed when SR95531 was applied after picrotoxin: ΔI_{hold} , -1.2 ± 0.9 pA, $n = 3$, $p = 0.3$). Subsequent application of picrotoxin resulted in an outward shift of I_{hold} by 18.9 ± 3.4 pA ($p = 0.0049$, $n = 5$), and a decrease in baseline RMS noise by 0.39 ± 0.05 pA ($p = 0.0012$, $n = 5$) in the same cells (Fig. 1B). This effect of picrotoxin was occluded by prior application of pentylenetetrazol (Table 1), another noncompetitive GABA_A antagonist (Huang et al., 2001). ΔI_{hold} following application of picrotoxin in the presence of pentylenetetrazol was 0.4 ± 0.7 pA ($n = 5$).

As previously reported (Semyanov et al., 2003), markedly increasing GABA in the perfusate to 5 μM revealed a SR95531-sensitive component of the tonic current (Fig. 1C). Under these conditions, application of a low concentration of SR95531 (0.5 μM) caused a significant outward shift of 29.4 ± 14.4 pA in the holding current ($p = 0.016$, $n = 8$) and a significant decrease in RMS noise by 3.0 ± 0.8 pA ($p = 0.008$, $n = 8$, Fig. 1C). Even in the presence of a high concentration of SR95531 (125 μM) under these conditions, application of picrotoxin caused a further outward shift in holding current by 9.2 ± 2.1 pA ($p = 0.031$, $n = 6$) and a reduction of RMS noise by 0.51 ± 0.09 pA ($p = 0.003$, $n = 6$; Fig. 1C).

The lack of efficacy of SR95531 under baseline conditions suggests that only negligible ambient GABA can be detected by dentate granule cells. Alternatively, SR95531 may not bind to the receptors mediating tonic current. The efficacy of SR95531 in expression systems in which $\alpha 4\delta$ subunit-containing receptors (the main contributor to tonic currents in dentate granule cells) (Glykys and Mody, 2007b) are expressed (Brown et al., 2002) argues against this. Nevertheless, we tested this using another competitive GABA_A antagonist, bicuculline. Application of bicuculline (10 μM) blocked all synaptic activity, induced a small outward shift of holding current by 5.9 ± 0.6 pA ($p = 0.00019$, $n = 6$), and decreased baseline RMS noise by 0.72 ± 0.11 pA ($p = 0.0014$, $n = 6$; Fig. 1D).

This result can be explained by the inverse agonist activity of bicuculline (Ueno et al., 1997; Bai et al., 2001; McCartney et al., 2007). If SR95531 does not bind to the same receptors as bicuculline, then it should have no effect in the presence of bicuculline. Conversely, if SR95531 does bind then it should displace the bicuculline and paradoxically induce an inward current. The effect of bicuculline was indeed reversed by application of SR95531 (25 μM), which activated an inward current of 10.7 ± 1.7 pA ($p = 0.0013$, $n = 6$), and increased baseline noise by 0.27 ± 0.06 pA ($p = 0.008$, $n = 6$). This was completely blocked by subsequent application of picrotoxin (Fig. 1D). These results suggest that all three GABA_A antagonists bind to the same receptors and that these receptors are not detecting GABA under baseline conditions. Moreover, the lack of effect of SR95531 in baseline conditions indicates that other possible endogenous agonists (e.g., taurine) are also not mediating the tonic current (Jia et al., 2008).

SR95531-insensitive tonic currents in DGCs are mediated by not only δ subunit-containing GABA_ARs

Tonic GABA_A-mediated currents in DGCs are predominantly mediated by $\alpha 4$ and δ subunit-containing receptors (Stell and Mody, 2002; Stell et al., 2003; Caraiscos et al., 2004). We therefore tested whether δ -GABA_ARs contribute to the SR95531-resistant tonic current. We took advantage of knock-out mice lacking δ subunits of GABA_ARs. In these mice, in contrast to rats, there was a very small SR95531-sensitive tonic current. However, the majority of the tonic current was SR95531 insensitive (Fig. 2). In $\delta^{-/-}$ mice in the presence of 25 μM SR95531, the effect of picrotoxin on holding current was $\sim 60\%$ less than that in wild-type mice (littermate controls), implying that δ -GABA_ARs contrib-

ute to SR95531-resistant tonic currents (Fig. 2). However, there was a significant SR95531-insensitive tonic current in $\delta^{-/-}$ mice, suggesting that other GABA_AR subtypes can also contribute to the SR95531-insensitive tonic current in DGCs (Glykys et al., 2008).

Low [GABA]_e is maintained both *in vitro* and *in vivo*

The fact that DGCs display SR95531-insensitive tonic currents suggests that [GABA]_e in *ex vivo* tissue is maintained at concentrations that are not detectable by high-affinity GABA_ARs.

We used GABA_AR openings in outside-out “sniffer” patches from dentate granule cells (Fig. 3A,B) to give a semiquantitative estimate of [GABA]_e in slices. We first confirmed that channel openings could be recorded in outside-out patches in the presence of 10 μ M GABA (channel openings disappeared in the absence of GABA) (Fig. 3A). These channels were blocked by picrotoxin and had an opening frequency of 20.1 ± 5.6 Hz, a conductance of 39.4 ± 7.2 pS, and an average open time of 32.1 ± 7.1 ms ($n = 11$). We were able to detect GABA_AR openings when the patch was 5 μ m above the slice (0.55 ± 0.05 Hz, $n = 5$), indicating that there is sufficient extracellular GABA to bind to GABA_ARs (Fig. 3B). These openings did not occur when the patch was moved 300 μ m above the slice and recurred on lowering the patch again to the slice (Fig. 3B). Consequent application of 200 nM GABA to the perfusate, comparable to that found in CSF (Glaeser and Hare, 1975), almost doubled the frequency of GABA_AR openings recorded above the slice surface. The effect of 200 nM GABA application was even more evident when patches were raised 300 μ m above the slice. This indicates that [GABA]_e is <200 nM in the slice (Fig. 3B) and is consistent with the predicted GABA transporter equilibrium [GABA]_e of ~ 100 nM (Wu et al., 2007). We further confirmed that similarly low [GABA]_e is maintained *in vivo* by using the zero-net-flux microdialysis method. Microdialysis probes were inserted into the CA3-dentate gyrus region via guide cannula, and *in vivo* [GABA]_e was calculated to be 92 ± 10 nM ($n = 5$; Fig. 3C,D).

Since an increase in GABA_AR openings occurs as the patch approaches the surface, yet there is no evidence of significant GABA-mediated tonic currents, it is likely that the activity of GABA transporters prevents extrasynaptic GABA_A receptors from being exposed to even such low concentrations under baseline condi-




Figure 3. Low extracellular GABA concentration *in vitro* and *in vivo*. **A**, Sniffer patches, outside-out patches from DGCs, display GABA_AR channel openings in the presence of GABA. Representative traces are shown on the left panel. Top, Application of 10 μ M GABA to the outside-out patches induces single-channel openings. Middle, A blow-up of the section of the top trace marked by the vertical dashed lines. Bottom, A trace illustrating no channel activity in the absence of GABA in perfusion solution. Right, Normalized all-points amplitude histogram from the experiment with 10 μ M GABA application. Horizontal lines indicate mean amplitude values of an open (O) and closed (C) state. **B**, The frequency of channel opening in the sniffer patch close to the slice surface (5 μ m above DGC layer) and in the bath (300 μ m above the slice surface) with and without 200 nM GABA in the perfusion solution (sample traces from the same patch at different conditions are shown on the left). Bar charts show (from top to bottom on the right) normalized values of channel opening frequency, open time, and conductance \pm SEM ($n = 5$; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$, paired Student's t test). **C**, Estimation of [GABA]_e in the rat hippocampus was performed using zero-net-flux microdialysis. Locations of the microdialysis probes in the hippocampus from five rats are shown by the black circles on the schematic of a horizontal section of rat brain (Paxinos and Watson, 2007). **D**, After the collection of six baseline samples, the microdialysis probe was perfused with increasing concentrations of GABA (C_{in} ; 10–240 nM) by reverse dialysis. The baseline condition represents $C_{in} = 0$. The concentration of GABA in the collected samples (C_{out} ; in nM) was determined by HPLC with electrochemical detection. The net loss or gain of GABA in the dialysate ($C_{out} - C_{in}$; in nM) was calculated and plotted against the concentration of GABA in the perfusion fluid (C_{in}). The intercept with the dashed line ($y = 0$) is the concentration of GABA at which $C_{out} = C_{in}$ represents an estimate of the [GABA]_e ($n = 5$).




Figure 4. Spontaneous GABA_AR channel openings in nucleated patches. **A**, Left, Representative traces illustrating single-channel openings; top to bottom: control, SR95531 (25 μM), picrotoxin (20 μM). Lower traces are expanded segments indicated by vertical dashed lines on the top traces. Right, Normalized all-points amplitude histograms. **B–D**, Channel opening frequency (**B**), conductance (**C**), and average open time (**D**) normalized to control values ($n = 5$ nucleated patches). **E**, Application of 125 μM SR95531 increases the frequency of spontaneous channel openings. Left, Sample traces. Right, Spontaneous channel opening frequency, open time, and conductance normalized to control values ($n = 5$). Horizontal dashed lines indicate an open (O) and closed (C) state. * $p < 0.05$; *** $p < 0.001$, paired t test. W/O, Washout.

tions. When GABA in the perfusate was increased to 200 nM, comparable to that found in CSF (Glaeser and Hare, 1975), I_{hold} did not change significantly and SR95531 (25 μM) had no significant effect on I_{hold} (ΔI_{hold} following consequent application of GABA: -1.21 ± 2.03 pA, $p = 0.6$; ΔI_{hold} following consequent application of SR95531: 1.7 ± 1.4 pA, $p = 0.29$; $n = 5$), indicating that there is no increase in the GABA detected by extrasynaptic GABA_ARs.

Spontaneously opening GABA_ARs in DGCs

The evidence thus far implies that tonic inhibition in DGCs *in situ* is predominantly GABA independent. Can tonic currents be mediated by spontaneously opening GABA_ARs? Such receptors have been reported in excised patch and cell-attached recordings from hippocampal neurons (Birnir et al., 2000a), but not in the DGCs (Birnir et al., 1994). Failure to detect spontaneously opening GABA_ARs may be due to the small area sampled using outside-




Figure 5. Application of GABA to nucleated patches from dentate granule cells induces SR95531-sensitive channel openings. **A**, Left, Representative traces illustrating single-channel openings at consecutive stages of the experiment performed in the presence of 10 μM GABA. From top to bottom: control, SR95531 (25 μM), washout of SR95531. Horizontal dashed lines indicate mean amplitude values of an open (O) and closed (C) state. Right, Normalized all-points amplitude histograms. **B–D**, Normalized average opening frequency (**B**), conductance (**C**), and open time (**D**) of channel openings induced by GABA application ($n = 5$ nucleated patches). ** $p < 0.01$, *** $p < 0.001$, paired Student's t test. W/O, Washout.

out patches and/or the dependence of such openings upon the internal neuronal environment. Indeed, we noted that in very few excised patches (<10%) could we detect infrequent channel openings (less than one every 20 s). We therefore performed recordings from nucleated patches (i.e., whole-cell excisions containing intact nuclei). In this preparation, spontaneous GABA_AR openings were recorded in the majority of patches (Fig. 4). These openings had a conductance (37.4 ± 6.1 pS) similar to that observed in excised outside-out patches with applied GABA (39 ± 7.2 pS; Fig. 3A), and comparable to that determined in other studies (Bright et al., 2011). As expected, spontaneous openings were not affected by SR95531 (25 μM) but were fully blocked by picrotoxin (Fig. 4A–D). Since a high concentration of SR 95521 (125 μM) had acted as a partial agonist (Fig. 1), we tested the effect of SR 95521 (125 μM) on spontaneous openings in nucleated patches and, as predicted, found that it increased the frequency of openings (Fig. 4E). Addition of 10 μM GABA to the nucleated patches revealed SR95531-sensitive channels with the same conductance (Fig. 5).

Discussion

Our results indicate that GABA-independent openings of GABA_ARs are the major contributor to tonic currents in dentate granule cells *ex vivo*, even when GABA concentrations in the

perfusate are increased to levels comparable to those measured *in vivo*.

We took advantage and confirmed the different pharmacologies of GABA_AR antagonists (Table 1), and have shown that *in situ* DGC tonic currents are resistant to SR95531, indicating that they are not being generated by GABA binding to the receptors. This is consistent with similar observations in cultured neurons, other brain areas and cell types (Birnie et al., 2000b; McCartney et al., 2007). Importantly, SR95531 has a paradoxical effect of inhibiting the action of the inverse agonist bicuculline. This result suggests that SR95531 is displacing bicuculline from the receptors and consequently that the lack of effect of SR95531 is not due to a failure to bind to the receptors that are mediating the tonic current. At high concentrations (125 μM), we also found that SR95531 can act as a partial agonist. A partial agonist effect of SR95531 (100 μM) and bicuculline (1 mM) has been described previously at GABA_ARs in which there is a point mutation in the γ subunit (Ueno et al., 1997). We did not test bicuculline at this concentration and so cannot exclude a partial agonist effect of very high concentrations of bicuculline at extrasynaptic receptors.

This lack of GABA binding explains why inhibiting vesicular GABA release in our studies has no effect on the tonic current. The low concentration of GABA despite vesicular release is al-

most certainly due to efficient GABA uptake, because, when GABA transporters are blocked, tonic currents become dependent upon vesicular GABA release (Glykys and Mody, 2007b). The effect of decreasing vesicular release has variable results depending upon the nature of the tissue studied, suggesting that GABA concentrations and the consequent detection of extracellular GABA by GABA_ARs vary between brain regions. For example, in thalamocortical neurons, 50 μ M SR95531 has been demonstrated to reveal robust tonic GABA_AR-mediated currents (Cope et al., 2005). There may even be local inhomogeneities of GABA concentrations due to regional differences in the distribution of GABA transporters (Semyanov et al., 2003). Indirect evidence of this in the hippocampus is supplied by the observation that tonic currents are reduced in CA1 interneurons, but not in pyramidal cells of GAD65-deficient mice (Song et al., 2011). However, whether the heterogeneity of extracellular GABA concentrations observed *in vitro* occurs *in vivo* is unclear.

Spontaneous openings of GABA_ARs have been previously described in some preparations (e.g., in isolated hippocampal pyramidal cells and in outside-out patches from pyramidal neurons, but not in DGCs). However, the role for these spontaneous openings, and whether they can contribute to GABA_AR-mediated signaling *in situ*, has remained unclear (Macdonald et al., 1989; Birnir et al., 2000b). Here, we recorded from nucleated patches to show directly that spontaneously opening receptors are present in dentate granule cells. As predicted from other studies (McCartney et al., 2007), they are not affected by the competitive GABA_AR antagonist SR95531 and are completely blocked by the open channel blocker picrotoxin. Lack of evidence of such openings in outside-out patches from dentate granule cells in previous studies may be related both to the small area of membrane and also to the effects of the internal cellular milieu on GABA_AR channel gating.

A vexed question is how GABA in a slice relates to the *in vivo* situation. Direct measurements of GABA in CSF have revealed concentrations on the order of 200 nM (Glaeser and Hare, 1975). Using the zero-net-flux microdialysis method, we have estimated the concentration in the extracellular fluid to be even lower. This is somewhat at odds with an earlier study that estimated the GABA concentration in the extracellular fluid of the hippocampus to be 800 nM (Lerma et al., 1986). This measurement, however, was confounded by the complex method used to retrieve absolute concentrations from dialysate concentrations and reliance on estimates of diffusion across the dialysis membrane. It should also be noted that these previous estimates were performed in rats under urethane anesthesia, and sampling was performed 1 h after probe implantation, when the blood–brain barrier could still be disrupted. The microdialysis method averages over a large area, and so we cannot exclude the possibility that the regional GABA concentrations may differ and, in particular, that the concentration detected by neurons may be greater than this depending upon the balance of local release and uptake; however, this was not observed by us (and others) using the sniffer patch technique. Indeed, 100 nM is close to the EC₂₀ for extrasynaptic δ -subunit-containing GABA_ARs (Bright et al., 2011), and so the absence of a detectable GABA-mediated current under baseline conditions or when GABA in the perfusate is increased to 200 nM indicates that the GABA detected by neurons is less than that present in the extracellular space, perhaps due to efficient local GABA uptake. These results strongly argue against the widespread addition of high GABA concentrations (usually 5 μ M) to the perfusate during *ex vivo* experiments. Indeed, these

concentrations of GABA are 50-fold greater than those we detect *in vivo*.

Our findings have several important implications. First, our results indicate that studies that use SR95531 to measure tonic currents may significantly underestimate them. The results may also explain discrepancies in the measurement of tonic currents in DGCs, which in one study (and in contrast to others) was proposed not to be present in control tissue because of the lack of effect of SR95531 (Zhan and Nadler, 2009). Furthermore, the widespread use of SR95531 and on occasion high concentrations of SR95531, which could have a partial agonist effect, may significantly underestimate the magnitude of tonic currents present. Even when GABA is gating the receptor, displacement by SR95531 would not prevent spontaneous openings, and may therefore not accurately measure the tonic current. Moreover, our results indicate that there is always a tonic current present in dentate granule cells even when extracellular GABA concentrations are low (as they are most of the time). The importance of maintaining such an inhibitory tone in neurons is further underscored by the observation that knocking out receptors generating tonic current in cerebellar granule cells leads to an upregulation of a two-pore potassium channel to maintain the conductance (Brickley et al., 2001).

References

- Attwell D, Barbour B, Szatkowski M (1993) Nonvesicular release of neurotransmitter. *Neuron* 11:401–407. [CrossRef Medline](#)
- Bai D, Zhu G, Pennefather P, Jackson MF, MacDonald JF, Orser BA (2001) Distinct functional and pharmacological properties of tonic and quantal inhibitory postsynaptic currents mediated by gamma-aminobutyric acid(A) receptors in hippocampal neurons. *Mol Pharmacol* 59:814–824. [Medline](#)
- Bennett MR, Kearns JL (2000) Statistics of transmitter release at nerve terminals. *Prog Neurobiol* 60:545–606. [CrossRef Medline](#)
- Birnir B, Everitt AB, Gage PW (1994) Characteristics of GABA_A channels in rat dentate gyrus. *J Membr Biol* 142:93–102. [Medline](#)
- Birnir B, Eghbali M, Everitt AB, Gage PW (2000a) Bicuculline, pentobarbital and diazepam modulate spontaneous GABA(A) channels in rat hippocampal neurons. *Br J Pharmacol* 131:695–704. [CrossRef Medline](#)
- Birnir B, Everitt AB, Lim MS, Gage PW (2000b) Spontaneously opening GABA(A) channels in CA1 pyramidal neurones of rat hippocampus. *J Membr Biol* 174:21–29. [CrossRef Medline](#)
- Brickley SG, Mody I (2012) Extrasynaptic GABA(A) receptors: their function in the CNS and implications for disease. *Neuron* 73:23–34. [CrossRef Medline](#)
- Brickley SG, Revilla V, Cull-Candy SG, Wisden W, Farrant M (2001) Adaptive regulation of neuronal excitability by a voltage-independent potassium conductance. *Nature* 409:88–92. [CrossRef Medline](#)
- Bright DP, Renzi M, Bartram J, McGee TP, MacKenzie G, Hosie AM, Farrant M, Brickley SG (2011) Profound desensitization by ambient GABA limits activation of delta-containing GABA_A receptors during spillover. *J Neurosci* 31:753–763. [CrossRef Medline](#)
- Brown N, Kerby J, Bonnert TP, Whiting PJ, Wafford KA (2002) Pharmacological characterization of a novel cell line expressing human α 4(3) β 2 GABA(A) receptors. *Br J Pharmacol* 136:965–974. [CrossRef Medline](#)
- Caraiscos VB, Elliott EM, You-Ten KE, Cheng VY, Belelli D, Newell JG, Jackson MF, Lambert JJ, Rosahl TW, Wafford KA, MacDonald JF, Orser BA (2004) Tonic inhibition in mouse hippocampal CA1 pyramidal neurons is mediated by α 5 subunit-containing gamma-aminobutyric acid type A receptors. *Proc Natl Acad Sci U S A* 101:3662–3667. [CrossRef Medline](#)
- Cope DW, Hughes SW, Crunelli V (2005) GABA_A receptor-mediated tonic inhibition in thalamic neurons. *J Neurosci* 25:11553–11563. [CrossRef Medline](#)
- Coulter DA, Carlson GC (2007) Functional regulation of the dentate gyrus by GABA-mediated inhibition. *Prog Brain Res* 163:235–243. [CrossRef Medline](#)
- Danglot L, Rostaing P, Triller A, Bessis A (2004) Morphologically identified

- glycinergic synapses in the hippocampus. *Mol Cell Neurosci* 27:394–403. [CrossRef Medline](#)
- de Groote L, Linthorst AC (2007) Exposure to novelty and forced swimming evoke stressor-dependent changes in extracellular GABA in the rat hippocampus. *Neuroscience* 148:794–805. [CrossRef Medline](#)
- Droste SK, de Groote L, Atkinson HC, Lightman SL, Reul JM, Linthorst AC (2008) Corticosterone levels in the brain show a distinct ultradian rhythm but a delayed response to forced swim stress. *Endocrinology* 149:3244–3253. [CrossRef Medline](#)
- Duveau V, Laustela S, Barth L, Gianolini F, Vogt KE, Keist R, Chandra D, Homanics GE, Rudolph U, Fritschy JM (2011) Spatiotemporal specificity of GABAA receptor-mediated regulation of adult hippocampal neurogenesis. *Eur J Neurosci* 34:362–373. [CrossRef Medline](#)
- Farrant M, Nusser Z (2005) Variations on an inhibitory theme: phasic and tonic activation of GABA(A) receptors. *Nat Rev Neurosci* 6:215–229. [CrossRef Medline](#)
- Gaspary HL, Wang W, Richerson GB (1998) Carrier-mediated GABA release activates GABA receptors on hippocampal neurons. *J Neurophysiol* 80:270–281. [Medline](#)
- Ge S, Goh EL, Sailor KA, Kitabatake Y, Ming GL, Song H (2006) GABA regulates synaptic integration of newly generated neurons in the adult brain. *Nature* 439:589–593. [CrossRef Medline](#)
- Glaeser BS, Hare TA (1975) Measurement of GABA in human cerebrospinal fluid. *Biochem Med* 12:274–282. [CrossRef Medline](#)
- Glykys J, Mody I (2007a) Activation of GABAA receptors: views from outside the synaptic cleft. *Neuron* 56:763–770. [CrossRef Medline](#)
- Glykys J, Mody I (2007b) The main source of ambient GABA responsible for tonic inhibition in the mouse hippocampus. *J Physiol* 582:1163–1178. [CrossRef Medline](#)
- Glykys J, Mann EO, Mody I (2008) Which GABA_A receptor subunits are necessary for tonic inhibition in the hippocampus? *J Neurosci* 28:1421–1426. [CrossRef Medline](#)
- Hamann M, Desarmenien M, Desaulles E, Bader MF, Feltz P (1988) Quantitative evaluation of the properties of a pyridazinyl GABA derivative (SR 95531) as a GABAA competitive antagonist. An electrophysiological approach. *Brain Res* 442:287–296. [CrossRef Medline](#)
- Herd MB, Haythornthwaite AR, Rosahl TW, Wafford KA, Homanics GE, Lambert JJ, Belelli D (2008) The expression of GABA_A beta subunit isoforms in synaptic and extrasynaptic receptor populations of mouse dentate gyrus granule cells. *J Physiol* 586:989–1004. [CrossRef Medline](#)
- Holter NI, Zylla MM, Zuber N, Bruehl C, Draguhn A (2010) Tonic GABAergic control of mouse dentate granule cells during postnatal development. *Eur J Neurosci* 32:1300–1309. [CrossRef Medline](#)
- Huang RQ, Bell-Horner CL, Dibas MI, Covey DF, Drewe JA, Dillon GH (2001) Pentylentetrazole-induced inhibition of recombinant gamma-aminobutyric acid type A (GABA(A)) receptors: mechanism and site of action. *J Pharmacol Exp Ther* 298:986–995. [Medline](#)
- Isaacson JS, Solis JM, Nicoll RA (1993) Local and diffuse synaptic actions of GABA in the hippocampus. *Neuron* 10:165–175. [CrossRef Medline](#)
- Jia F, Yue M, Chandra D, Keramidias A, Goldstein PA, Homanics GE, Harrison NL (2008) Taurine is a potent activator of extrasynaptic GABA_A receptors in the thalamus. *J Neurosci* 28:106–115. [CrossRef Medline](#)
- Jones MV, Jonas P, Sahara Y, Westbrook GL (2001) Microscopic kinetics and energetics distinguish GABA(A) receptor agonists from antagonists. *Biophys J* 81:2660–2670. [CrossRef Medline](#)
- Kuntz A, Clement HW, Lehnert W, van Calker D, Hennighausen K, Gerlach M, Schulz E (2004) Effects of secretin on extracellular amino acid concentrations in rat hippocampus. *J Neural Transm* 111:931–939. [CrossRef Medline](#)
- Lee S, Yoon BE, Berglund K, Oh SJ, Park H, Shin HS, Augustine GJ, Lee CJ (2010) Channel-mediated tonic GABA release from glia. *Science* 330:790–796. [CrossRef Medline](#)
- Lerma J, Herranz AS, Herreras O, Abaira V, Martín del Río R (1986) In vivo determination of extracellular concentration of amino acids in the rat hippocampus. A method based on brain dialysis and computerized analysis. *Brain Res* 384:145–155. [CrossRef Medline](#)
- Linthorst AC, Flachskamm C, Holsboer F, Reul JM (1994) Local administration of recombinant human interleukin-1 beta in the rat hippocampus increases serotonergic neurotransmission, hypothalamic-pituitary-adrenocortical axis activity, and body temperature. *Endocrinology* 135:520–532. [CrossRef Medline](#)
- Macdonald RL, Rogers CJ, Twyman RE (1989) Kinetic properties of the GABAA receptor main conductance state of mouse spinal cord neurones in culture. *J Physiol* 410:479–499. [Medline](#)
- Mann EO, Mody I (2010) Control of hippocampal gamma oscillation frequency by tonic inhibition and excitation of interneurons. *Nat Neurosci* 13:205–212. [CrossRef Medline](#)
- Martin LJ, Zurek AA, MacDonald JF, Roder JC, Jackson MF, Orser BA (2010) $\alpha 5$ GABA_A receptor activity sets the threshold for long-term potentiation and constrains hippocampus-dependent memory. *J Neurosci* 30:5269–5282. [CrossRef Medline](#)
- McCartney MR, Deeb TZ, Henderson TN, Hales TG (2007) Tonic active GABAA receptors in hippocampal pyramidal neurons exhibit constitutive GABA-independent gating. *Mol Pharmacol* 71:539–548. [CrossRef Medline](#)
- Mihalek RM, Banerjee PK, Korpi ER, Quinlan JJ, Firestone LL, Mi ZP, Laganaur C, Tretter V, Sieghart W, Anagnostaras SG, Sage JR, Fanselow MS, Guidotti A, Spiegelman I, Li Z, DeLorey TM, Olsen RW, Homanics GE (1999) Attenuated sensitivity to neuroactive steroids in γ -aminobutyrate type A receptor δ subunit knockout mice. *Proc Natl Acad Sci U S A* 96:12905–12910. [CrossRef Medline](#)
- Mtchedlishvili Z, Kapur J (2006) High-affinity, slowly desensitizing GABAA receptors mediate tonic inhibition in hippocampal dentate granule cells. *Mol Pharmacol* 69:564–575. [CrossRef Medline](#)
- Mtchedlishvili Z, Lepsveridze E, Xu H, Kharlamov EA, Lu B, Kelly KM (2010) Increase of GABAA receptor-mediated tonic inhibition in dentate granule cells after traumatic brain injury. *Neurobiol Dis* 38:464–475. [CrossRef Medline](#)
- Nusser Z, Mody I (2002) Selective modulation of tonic and phasic inhibitions in dentate gyrus granule cells. *J Neurophysiol* 87:2624–2628. [Medline](#)
- Nyitrai G, Kékesi KA, Juhász G (2006) Extracellular level of GABA and Glu: in vivo microdialysis-HPLC measurements. *Curr Top Med Chem* 6:935–940. [CrossRef Medline](#)
- Overstreet LS, Westbrook GL (2001) Paradoxical reduction of synaptic inhibition by vigabatrin. *J Neurophysiol* 86:596–603. [Medline](#)
- Pavlov I, Savtchenko LP, Kullmann DM, Semyanov A, Walker MC (2009) Outwardly rectifying tonically active GABA_A receptors in pyramidal cells modulate neuronal offset, not gain. *J Neurosci* 29:15341–15350. [CrossRef Medline](#)
- Paxinos G, Watson C (2007) The rat brain in stereotaxic coordinates, Ed 6. London: Elsevier.
- Rossi DJ, Hamann M, Attwell D (2003) Multiple modes of GABAergic inhibition of rat cerebellar granule cells. *J Physiol* 548:97–110. [CrossRef Medline](#)
- Semyanov A, Walker MC, Kullmann DM (2003) GABA uptake regulates cortical excitability via cell type-specific tonic inhibition. *Nat Neurosci* 6:484–490. [CrossRef Medline](#)
- Semyanov A, Walker MC, Kullmann DM, Silver RA (2004) Tonic active GABA A receptors: modulating gain and maintaining the tone. *Trends Neurosci* 27:262–269. [CrossRef Medline](#)
- Song I, Savtchenko L, Semyanov A (2011) Tonic excitation or inhibition is set by GABA(A) conductance in hippocampal interneurons. *Nat Commun* 2:376. [CrossRef Medline](#)
- Stell BM, Mody I (2002) Receptors with different affinities mediate phasic and tonic GABA_A conductances in hippocampal neurons. *J Neurosci* 22:RC223. [Medline](#)
- Stell BM, Brickley SG, Tang CY, Farrant M, Mody I (2003) Neuroactive steroids reduce neuronal excitability by selectively enhancing tonic inhibition mediated by delta subunit-containing GABAA receptors. *Proc Natl Acad Sci U S A* 100:14439–14444. [CrossRef Medline](#)
- Sylantyev S, Savtchenko LP, Niu YP, Ivanov AI, Jensen TP, Kullmann DM, Xiao MY, Rusakov DA (2008) Electric fields due to synaptic currents sharpen excitatory transmission. *Science* 319:1845–1849. [CrossRef Medline](#)
- Traynelis SF, Jaramillo F (1998) Getting the most out of noise in the central nervous system. *Trends Neurosci* 21:137–145. [CrossRef Medline](#)
- Ueno S, Bracamontes J, Zorumski C, Weiss DS, Steinbach JH (1997) Bicuculline and gabazine are allosteric inhibitors of channel opening of the GABA_A receptor. *J Neurosci* 17:625–634. [Medline](#)
- Wu Y, Wang W, Diez-Sampedro A, Richerson GB (2007) Nonvesicular inhibitory neurotransmission via reversal of the GABA transporter GAT-1. *Neuron* 56:851–865. [CrossRef Medline](#)
- Yang Z, Cromer BA, Harvey RJ, Parker MW, Lynch JW (2007) A proposed

structural basis for picrotoxinin and picrotin binding in the glycine receptor pore. *J Neurochem* 103:580–589. [CrossRef Medline](#)
Zhan RZ, Nadler JV (2009) Enhanced tonic GABA current in normotopic

and hilar ectopic dentate granule cells after pilocarpine-induced status epilepticus. *J Neurochem* 102:670–681.